



REMOTE ISCHEMIC PRECONDITIONING IS NOT MEDIATED BY ENDOGENOUS BRADYKININ IN HUMANS

ACC Oral Contributions

Ernest N. Morial Convention Center, Room 228

Monday, April 04, 2011, 5:15 p.m.-5:30 p.m.

Session Title: Endothelial Function and Atherosclerotic Vascular Disease

Abstract Category: 8. Vascular Biology/Atherosclerosis/Thrombosis/Endothelium

Presentation Number: 919-5

Authors: Christian M. Pedersen, Gareth Barnes, Michael R. Schmidt, Hans Erik Botker, Rajesh K. Kharbada, David E. Newby, Nicholas LM Cruden, University of Edinburgh, Centre for Cardiovascular Science, Edinburgh, United Kingdom, Aarhus University Hospital Skejby, Cardiology Research, Aarhus, Denmark

Background Ischemia reperfusion injury (IRI) induces endothelial dysfunction with attenuation of acetylcholine-induced vasodilatation in man. Remote ischemic preconditioning (RIPC) preserves endothelium-dependent vasodilatation following IRI. Given that exogenous bradykinin protects against IRI, we hypothesised that endogenous bradykinin mediates the vascular protective effects of RIPC.

Methods In a double blind crossover study, 20 subjects were randomised to receive an intravenous infusion of the bradykinin B2 receptor antagonist, HOE-140 (0.1 mg/kg body weight) or saline placebo. IRI (20 min upper arm cuff inflation to 200 mmHg) was induced in the non-dominant arm in all subjects. Prior to induction of IRI 10 subjects also received 3 cycles of 5 min RIPC in the dominant arm. Using bilateral forearm venous occlusion plethysmography, blood flow was measured during intra-brachial infusion of acetylcholine (ACh; 5-20 µg/min) at baseline and following IRI.

Results Compared to baseline, acetylcholine-induced vasodilatation was reduced at 15 and 45 min following IRI, both in the presence ($P=0.0002$) and absence ($P=0.04$) of HOE-140. RIPC prevented the reduction in ACh-induced vasodilatation associated with IRI, irrespective of the presence or absence of HOE-140 ($P=NS$ for both).

Conclusions Endothelial vasomotor dysfunction is induced by IRI and can be prevented by RIPC. Our findings suggest that endogenous bradykinin does not play a major role in the induction of IRI or its prevention by RIPC.

